



COMSATS University, Islamabad (CUI)  
Islamabad Campus

Fall 2024  
Midterm Exam

Class: BCS-R

Subject: Multivariable Calculus (MTH 105)


Instructor: Dr. Manshoor Ahmed

Date: 05-11-2024, Time [11:30-12:50]

Q1. (i) Which of the following is the equation of straight line  $x = a$  in polar coordinates:

(a)  $r = a \cos \theta$  (b)  $r = a \sin \theta$  (c)  $r = a \sec \theta$  (d)  $r = a \csc \theta$ .

(ii) The equation of the given cylindrical surface



(a)  $y = x^2$  (b)  $z = \cos x$  (c)  $z = x^2$  (d)  $z = \sin x$

(iii) Which of the following point lie on the curve  $r = a \cos \theta$  is

(a)  $(-a, \pi)$ . (b)  $(a, \pi)$  (c)  $(a, \pi/2)$ . (d)  $(a, 0)$

(iv) If  $f(x, y) = \sin(xy) + x^2 \ln(y)$ , then  $f_{yx}$  at  $(0, \pi/2)$ :

(a) 33 (b) 0 (c) 3 (d) 1

(v) The function  $f(x, y) = x^2 + 2xy - y^3$

(a) Only one stationary point at  $(0,0)$ . (b) Two stationary point at  $(0,0)$  and  $(1/6, 1/3)$ .

(c) Two stationary point at  $(0,0)$  and  $(1/6, 2/3)$ . (d) No stationary point.

Q2. Sketch and find the area of the region that is outside the cardioid  $r = 2 - 2 \cos \theta$  and inside the circle  $r = 4$ .

Q3. Find the eccentricity and the distance from the pole to the directrix and sketch the plot in polar coordinates.

$$r = \frac{4}{3 - 2 \cos \theta}.$$

Q4. Solve the following problem by using Lagrange Multipliers. Find the point on the circle  $x^2 + y^2 = 45$  that are closest and farthest from  $(1,2)$ .

Q#2

Points of intersection  
of two curves.

$$r = 2 - 2\cos\theta \quad \& \quad r = 4$$

$$2 - 2\cos\theta = 4$$

$$1 - \cos\theta = 2$$

$$\cos\theta = -1$$

$$\Rightarrow \theta = \cos^{-1}(-1) = \pi$$

$\therefore$  Required area.

$$A = 2 \left[ \frac{1}{2} \int_{\pi}^{\pi} \left[ (4)^2 - (2 - 2\cos\theta)^2 \right] d\theta \right]$$

$2 \times \frac{1}{2}$  due to symmetry.

$$A = \int_0^{\pi} 16 d\theta - \int_0^{\pi} (4 + 4\cos^2\theta - 8\cos\theta) d\theta$$

$$= 16 \left[ \theta \right]_0^{\pi} - 4 \int_0^{\pi} (1 + \cos^2\theta - 2\cos\theta) d\theta$$

$$= 16\pi - 4 \int_0^{\pi} \left( 1 + \left( \frac{1 + \cos 2\theta}{2} \right) - 2\cos\theta \right) d\theta$$

$$= 16\pi - 4\pi - \frac{4}{2} \left[ \theta + \frac{\sin 2\theta}{2} \right]_0^{\pi} - 8 \left[ \sin\theta \right]_0^{\pi}$$

$$= 12\pi - 2\pi - 0 - 8(0-0)$$

$$= 8\pi$$

x ————— y

Q#3


Given equation is

$$r = \frac{4}{3 - 2\cos\theta}$$

$$r = \frac{4/3}{1 - \frac{2}{3}\cos\theta} = \frac{\frac{2}{3}(2)}{1 - \frac{2}{3}\cos\theta}$$


Here,  $e = \frac{2}{3}$ ,  $d = 2$ , Further  $0 < e < 1$ .

$\therefore$  the given eq. represents ellipse.



## Table of value.

|          |   |         |       |          |        |
|----------|---|---------|-------|----------|--------|
| $\theta$ | 0 | $\pi/2$ | $\pi$ | $3\pi/2$ | $2\pi$ |
| $r$      | 4 | $4/3$   | $4/5$ | $4/3$    | 4      |



$$\theta = 0$$

$$r = \frac{4}{3 - 2\cos 0} = 4$$

$$\theta = \pi/2$$

$$r = \frac{4}{3 - 2\cos \pi/2} = 4/3$$

$$\theta = \pi$$

$$r = \frac{4}{3 - 2\cos \pi} = 4/5$$

$$\theta = 3\pi/2$$

$$r = \frac{4}{3 - 2\cos 3\pi/2} = 4/3$$

$$\theta = 2\pi$$

$$r = \frac{4}{3 - \cos 2\pi} = 4$$

Q #4 Let  $(x, y)$  be the point on the given circle,

then distance of  $(x, y)$  from  $(1, 2)$  is

$$f(x, y) = D^2 = (x-1)^2 + (y-2)^2$$

$$f(x, y) = (x-1)^2 + (y-2)^2$$

Subject to  $g(x, y) = x^2 + y^2 - 45 \rightarrow (1)$

$$\nabla f(x, y) = 2(x-1)\hat{i} + 2(y-2)\hat{j} \quad \nabla g = 2x\hat{i} + 2y\hat{j}$$

Since  $\nabla g \neq 0$  at the circle.  $\therefore$  Extreme values exist at the points, where,

$$\nabla f = \lambda \nabla g$$

$$\Rightarrow 2(x-1)\hat{i} + 2(y-2)\hat{j} = \lambda(2x\hat{i} + 2y\hat{j})$$

Comparison gives

$$2(x-1) = \lambda(2x), \quad 2(y-2) = \lambda(2y)$$

$$\Rightarrow \frac{x-1}{x} = \lambda, \quad \frac{y-2}{y} = \lambda$$

$$\frac{x-1}{x} = \frac{y-2}{y} \Rightarrow y(x-1) = x(y-2)$$

$$xy - y = xy - 2x$$

$$y = 2x \rightarrow (ii)$$

Using (ii) into (i)

$$x^2 + (2x)^2 = 45 \Rightarrow x^2 = 9 \Rightarrow x = \pm 3$$

When  $x = +3$  then  $y = 6$

When  $x = -3$  then  $y = -6$ .

$\therefore$  possible points are.

$$(3, 6), (-3, -6)$$

$$\text{As } f(3, 6) = (3-1)^2 + (6-2)^2 = 20$$

$$f(-3, -6) = (-3-1)^2 + (-6-2)^2 = 20.$$

Therefore,  $(3, 6)$  is closet and  $(-3, -6)$  is farthest.

$\lambda$  —————  $x$

